

Review paper

Changing Pattern of Mechanical Bowel Obstruction and Management Outcome in North-Eastern Nigeria

S Aliyu*, MB Tahir, UD Babayo, AB Zarami, AG Ibrahim and AG Madziga

Department of Surgery and Histopathology, University of Maiduguri Teaching Hospital Maiduguri, Borno State, Nigeria.

Accepted 23rd, September 2017

Abstract

Background: The study reviewed mechanical bowel obstruction over a five year period in North-Eastern Nigeria. **Patients and Methods:** The study retrospectively reviewed all patients that presented with mechanical bowel obstruction in a District Hospital Damaturu North –Eastern Nigeria managed between January 2011 and December 2015. **Results:** A total of 94 patients were managed age ranged between 20 and 72 years with male to female ratio of 1.4:1. The peak age group was 21- 30years accounting for 29.78%. Tumour was the commonest cause in 27.66% followed by external hernias and intra peritoneal adhesions 25.53% each. The procedures carried were bowel resection in 54.26% and herniorrhapy in 25.53%. The post operative complications were surgical site infection in 22.34%, enterocutaneous fistula in 3.19%. The mortality recorded was 15.96% majority due to metastatic colonic tumour. **Conclusion:** The rising incidence of colonic tumours and late presentation in the developing world and falling complicated external hernias due to availability of elective operations makes the former to become the most frequent cause of mechanical bowel obstruction in this environment.

Keywords: Mechanical bowel obstruction, Changing pattern, Management outcome.

INTRODUCTION

Mechanical bowel obstruction (MBO) is a common surgical emergency and frequently encountered problem in abdominal surgery. It constitutes a major cause of morbidity and a significant cause of surgical emergency. It requires a quick diagnosis as well as immediate rational and effective treatment. The surgeons are concerned with it because strangulation, causing bowel ischemia, necrosis and perforation might be involved, and it is often difficult to distinguish simple obstruction from strangulation obstruction. Accurate early diagnosis of intestinal strangulation in patients with mechanical bowel obstruction is important to decide on emergency surgery or to allow safe non operative management of carefully selected patients. Mechanical bowel obstruction is one of the commonest indications for emergency laparotomy,

perhaps second to peritonitis. Among such causes of mechanical obstructions are obstructed external hernias, volvulus, colonic tumour obstruction, and adhesions, Andrew, 2006; Daniel and Zuri, 2012. There is a changing pattern in clinical presentation of external abdominal hernias favouring elective procedures for such hernias when compared to a decade ago where complicated hernia presentation were common. There is a rising incidence in colonic tumours with bowel obstruction at presentation, Micheil and Agis, 2016. The rising incidence is being attributed to changing diet, urbanization with its attendant social habit like alcohol ingestion and smoking, Marinos *et al.*, 2013; Su and Arab, 2004; David *et al.*, 2010. One of the cardinal principles in the management of intestinal obstruction is adequate resuscitation before definitive surgery in order to reduce morbidity and mortality, Zheng-shui *et al.*, 2017. The aim of this study was to determine the pattern and outcome of mechanical bowel obstruction.

PATIENTS AND METHODS

The study retrospectively reviewed all patients presented with mechanical bowel obstruction between January 2011 and December 2015. Permission for the study was granted by the Hospital management and informed consent obtained from all patients. Information extracted from clinical and laboratory records and data analyzed using SPSS statistical analysis. All patients were resuscitated using intravenous fluids, antibiotics (ceftriaxone/ metronidazole), tetanus toxoid, blood, and diverting colostomy where necessary. Investigations done were full blood count; blood chemistry, random blood sugar, proctosigmoidoscopy, colonoscopy and biopsy. Others were barium enema, abdominopelvic ultrasound scan; chest x-ray, plain abdominal x-rays (erect and supine), and ECG. Computerized Tomography scan and MRI were done where indicated. All patients had bowel preparation before definitive surgery under general anesthesia.

RESULTS

A total of 94 patients were managed age ranged between 20 and 72 years with male to female ratio of 1.4:1. The peak age group was 21- 30years accounting for 28(29.78%) table 1. Tumour was the commonest cause in 26(27.66%) table 2. The procedures carried were bowel resection in 51(54.26%) table 3. The post operative complications were surgical site infection in 21(22.34%), and enterocutaneous fistula in 3(3.19%), which closed on conservative treatment. The mortality recorded was 7(7.45%), and 5(5.32%) due to metastatic colonic tumour.

DISCUSSION

The incidence of MBO was found to be higher among the age groups 21-30 years accounting for 29.78% which is similar to the findings by Soressa *et al.*, (2016). Colonic tumours was found to be the commonest cause of mechanical bowel obstruction in 44.68% which is in sharp contrast to similar study in the same environment a decade ago by Madziga *et al.*, 2008, that reported external hernias as the commonest. Obstructed external hernias are declining due to increase in elective herniorrhapy on the other hand there is increasing incidence in colonic tumours in the young due to change in diet and social habit. The highest procedure performed was bowel resection because it is the recommended primary treatment for volvulus and tumour as a global best practice, Osiro *et al.*, 2012; Sedat *et al.*, 2014. In addition, late strangulation of hernia contributed to the high bowel resection rate. The commonest post operative complication was surgical site infection in 22.34% which

was similar to the findings by Chang *et al.*, 2000, due to the fact that colonic procedures are dirty. Enterocutaneous fistula was 3.19%, due to the fact that colonic anastomosis has potential to leak. This was similar to the findings by Moghadamyeghaneh *et al.*, 2016, that recorded 3.8% in their studies. The current study had a mortality of 7%, which was higher than Bjorg *et al.*, 2016, who recorded a declining mortality of 4% from 5% over a decade. The higher mortality in this study can be attributed to tumour patients presenting with metastatic disease.

TABLE 1: Age Distribution

Age years	No	%
10-20	2	2.13
21-30	28	29.78
31-40	12	12.77
41-50	16	17.02
51-60	26	27.66
61-70	8	8.51
71-80	2	2.13
Total	94	100.00

TABLE 2: Diagnosis

Diagnosis	No	%
Hernia	24	25.53
Tumour	42	44.68
Volvulus	4	4.24
Adhesion	24	25.53
Faecal mass	3	3.19
Total	94	100.00

TABLE 3: Procedures

Procedures	NO
Herniorrhapy	24
Bowel resection	51
Adhesiolysis	24
Faecal evacuation	3

NB: Bowel resection: Tumour- 42, Volvulus-4, hernia-5

REFERENCES

Andrew K. The management of Incisional Hernia. Ann R Coll Surg. Engl. 2006 may; 88(3): 252 – 260
 Bjorg T.F., Jonas F., Lodve S., Odd S., Knut S., Asgaut V. Complications and death after Surgicval Treatment of Small Bowel Obstruction. Ann Surg. 2000 Apr; 231(4):529=537
 Chang H Hall GA Geerts WH Greenwood Cmckleod RSSher GD Allogeneircd blood cell transfusion is an independent risk factor for the development of postoperative bacterial infection. Vox

- sang 2000; 7813-18
- Daniel G., Zuri M. Management of colonic volvulus. *Clin Colon Rectal Surg.* 2012 Dec; 25(4): 236 – 244
- David L., Robert A.V., Lori S. T., Alice H. W. Cigarette Smoking and Colorectal cancer Risk by Molecularly Defined Subtypes. *J Natl Cancer Inst.* 2010 July 21; 102(14):1012- 1022
- Evaghelos X., Nicolaos G., Charina T. Et al. Clinical practice guidelines for the surgical management of colon cancer: a consensus statement of the Hellenic and Cypriot colorectal cancer study Group by the HeSMO. *Ann Gastroenterol .* 2016 Jan-March; 29(1): 3 – 17
- Lewis S. P. Laparotomy for Intestinal obstruction. *Ann Surg.* 1888, 3(3): 214 – 221
- Madziga A.G, Nuhu A. Causes and treatment outcome of mechanical bowel obstruction in north eastern Nigeria. *West Afri J Med.* 2008 Apr; 27(2):101 – 5
- Marinos P., Dalvinder M., Martyn E. C. Diet and supplements and their impact on colorectal cancer. *J. Gastrointest Oncol* 2013 Dec; 4(4): 409 – 423
- Micheil K., Agis E. Increasing incidence of colon cancer in patients< 50 years old: a new entity? *Ann Transl Med.* 2016 May; 4(9): 164 doi:10.21037/atm.2016.04.13
- Moghadamyeghanneh Z., Hannah M.H., Alizadeh R.F., Carmichael J.C., Mills S., Pigazzi A., Styamos M.J. Contemporary Management of anastomotic leak after colon surgery: assessing the need for reoperation. *Am J Surg.* 2016 Jun; 211(6):1005-13
- Osiro S.B., Cunningham D., Tubbs R.S., Gielecki J., Loukas M. The twisted colon: a review of sigmoid volvulus. *Am Surg.* 2012 Mar; 78(3): 271-9
- Sedat B., Huseyin O.A., Erdal K., Hakan Y., Fazilet K., Sedat Y. Outcomes of Surgical Treatment of Primary Signet Ring Cell Carcinoma of the Colon and Rectum: 22 Cases Reviewed With Literature. *Int Surg.* 2014Nov-Dec; 99(6):691-698
- Su L.J., Arab L. Alcohol consumption and risk of colon cancer: evidence from the national health and nutrition examination survey 1 epidemiological follow-up study. *Nutr cancer.* 2004; 50(2): 111 – 9
- Urgessa S., Abebe M., Desta H., Netsanet F. Prevalence, causes and management outcome of intestinal obstructyion in Adama Hospital, Ethiopia. *BMC Surgery* (2016) 16:38, DOI10.1186/s12893-0150-5
- Zheng-shui X., Wei X., Jia-qi Y., Hua C. Mechanical intestinal obstruction Secondary to Appendiceal mucinous cystadenoma. *Medicine (Baltimore).* 2017 feb; 95(5): e6016